

**Title:** Cost-Effectiveness of Transcatheter versus Surgical Aortic Valve Replacement in Patients at Lower Surgical Risk: Results from the NOTION Trial.

**Authors:** Benjamin P. Geisler, M.D; Troels H. Jørgensen, M.D; Hans Gustav H. Thyregod, M.D, PhD; Jan B. Pietzsch, PhD; Lars Søndergaard, M.D, PhD

**DOI:** 10.4244/EIJ-D-18-00847

**Citation:** Geisler BP, Jørgensen TH, Thyregod HGH, Pietzsch JB, Søndergaard L. Cost-Effectiveness of Transcatheter versus Surgical Aortic Valve Replacement in Patients at Lower Surgical Risk: Results from the NOTION Trial. *EuroIntervention* 2019; Jaa-632 2019, doi: 10.4244/EIJ-D-18-00847

**Guest Editor:** Alec Vahanian, M.D, PhD

**Manuscript submission date:** 20 August 2018

**Revisions received:** 08 April 2019, 28 May 2019, 01 July 2019, 15 July 2019

**Accepted date:** 13 August 2019

**Online publication date:** 20 August 2019

**Disclaimer:** This is a PDF file of a "Just accepted article". This PDF has been published online early without copy editing/typesetting as a service to the Journal's readership (having early access to this data). Copy editing/typesetting will commence shortly. Unforeseen errors may arise during the proofing process and as such Europa Digital & Publishing exercise their legal rights concerning these potential circumstances.

# **Cost-Effectiveness of Transcatheter versus Surgical Aortic Valve Replacement in Patients at Lower Surgical Risk**

Results from the NOTION Trial

Benjamin P. Geisler, M.D.<sup>1,2</sup>, Troels H. Jørgensen<sup>3</sup>, Hans Gustav H. Thyregod, M.D., Ph.D.<sup>3</sup>, Jan B. Pietzsch, Ph.D.<sup>1</sup>, and Lars Søndergaard, M.D., Ph.D.<sup>3</sup>

<sup>1</sup>Wing Tech Inc., Menlo Park, CA, USA; <sup>2</sup>Department of Medicine, Massachusetts General Hospital/Harvard Medical School, Boston, MA, USA; <sup>3</sup>The Heart Center, Rigshospitalet, Copenhagen, Denmark

**Short Running title:** Cost-effectiveness of TAVI in Lower Risk Patients

**Corresponding author:** Jan B. Pietzsch, PhD, Wing Tech Inc., 101 Jefferson Drive, Menlo Park, CA 94025, USA; [jpietzsch@wing-tech.com](mailto:jpietzsch@wing-tech.com)

**Conflict of Interest Statement/Funding:** Wing Tech Inc. (J.B.P., B.P.G.) provided health-economic consulting services to Medtronic plc. L.S. was the principal investigator for the NOTION trial and has received consultant fee and institutional research grants from Medtronic plc. H.G.H.T. was the other principal investigator for the NOTION trial and has no conflicts of interest to declare. ICMJE statements are attached.

*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*

## ABSTRACT

**Aims:** To estimate the cost-effectiveness of transcatheter aortic valve implantation (TAVI) vs surgical aortic valve replacement (SAVR) in patients at lower surgical risk.

**Methods and results:** Discounted costs from a societal perspective and effectiveness as quality-adjusted life years (QALYs) were projected to lifetime via a decision-analytic model calibrated to 60-month data from the NOTION trial. The base case assumed a scenario in which any mortality benefit would gradually fade out over time, with other scenarios explored in sensitivity analyses. The incremental cost-effectiveness ratio (ICER) was compared to the country-specific willingness-to-pay (WTP) threshold of 1.13 million Danish kroner (DKK). The base case ICER was DKK696,264/QALY (around €72,100/QALY via purchasing parity adjustment). Variation in long-term mortality beyond 5 years led to limited variation of incremental costs (DKK64,200 to 64,600), but a more pronounced variation in incremental QALYs (0.07 to 0.19 QALYs for most conservative and optimistic assumptions, compared to base case of 0.09 QALYs). All resulting ICERs (range DKK334,200 to DKK904,100 per QALY gained) were below the WTP threshold.

**Conclusions:** TAVI in a cohort of primarily low-surgical-risk patients was found to be a cost-effective treatment strategy in the Danish healthcare system. Cost-effectiveness analyses in other settings are warranted as are registries given the sensitivity of the model to long-term mortality.

## **Classifications**

TAVI; Death; No specific risk; Miscellaneous

## **Condensed Abstract**

Transcatheter aortic valve implantation (TAVI) provides outcomes similar to those of surgical aortic valve replacement (SAVR) in patients with severe aortic stenosis at lower surgical risk, but the cost-effectiveness in this patient group remains unknown. A decision-analytic model projected discounted quality-adjusted life time and costs based on the 60-month results of the NOTION trial. The base case incremental cost effectiveness ratio was DKK696,264/QALY. Long-term mortality influenced the results. All scenarios were below a country-specific willingness-to-pay threshold, rendering TAVI – compared to SAVR – a cost-effective treatment strategy in this Danish cohort of patients at lower surgical risk.

## **Abbreviations**

DKK=Danish krone

ICER=incremental cost-effectiveness ratio

MI=myocardial infarction

NOTION=Nordic Aortic Valve Intervention Trial

NYHA=New York Heart Association

PPM=permanent pacemaker implantation

PPP=purchasing power parity

QALY=quality-adjusted life year

*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*

RR=relative risk

SAVR=surgical aortic valve replacement

TAVI=transcatheter aortic valve implantation

## INTRODUCTION

Transcatheter aortic valve implantation (TAVI) has experienced a remarkable uptake in patients with severe aortic stenosis and high surgical risk, with over 54,000 procedures between 2012 and 2015 in the United States alone <sup>1</sup>. Its cost-effectiveness in these populations has been established in various settings and for both balloon-expandable and self-expandable TAVI devices <sup>2</sup>.

The Nordic Aortic Valve Intervention (NOTION) trial was the first trial in patients with severe aortic stenosis and lower surgical risk randomized to TAVI or surgical bioprosthetic aortic valve replacement (SAVR) <sup>3,4</sup>; 4-year results were recently presented <sup>5</sup>. The NOTION trial demonstrated no significant difference in the primary outcome, a composite of all-cause death, stroke, or myocardial infarction (MI), between TAVI and SAVR. Non-significant differences persisted after 24 and 48 months of follow-up, respectively <sup>4,5</sup>.

Given that surgical candidates who are not at increased surgical risk might have fewer (or no) incremental benefits from TAVI compared to SAVR but potentially higher costs, the “real-world” value of TAVI outside carefully selected high or extreme surgical risk patient cohorts has been uncertain. The objective of the present study was therefore to estimate the long-term cost-effectiveness of TAVI compared to SAVR in the NOTION cohort.

## **METHODS**

A Markov state transition model, nested in a decision tree (see Figure 1), was developed to compute long-term costs and effectiveness of TAVI vs. SAVR from the NOTION trial, as well as to quantify the impact of variation in the following input parameters: trial-based efficacy and adverse events, procedure and event costs, and New York Heart Association class (NYHA)-stratified health-related quality of life estimates scored for the Danish population. Patient-level data from the 6-year follow-up were analyzed wherever possible. The study adheres to both the American College of Cardiology/American Heart Association Statement on Cost/Value Methodology in Clinical Practice Guidelines and Performance Measures as well as the Consolidated Health Economic Evaluation Reporting Standards statement<sup>6,7</sup>. Please see Table 1 and Supplementary Tables 1-5 for the input parameters. The complete methods are described in the Supplementary Text.

## **RESULTS**

### **Model Calibration and Base Case Findings**

After calibrating the available 60 months of data, the model output matched the actual trial results for the endpoints all-cause mortality, stroke, and MI; see the Supplementary Figures 1-3 for details.

The model projected life expectancy was 8.95 and 8.76 life years for TAVI and SAVR, respectively, a difference of 69 days. Consequently, TAVI and SAVR patients would have lived until 88.0 and 87.9 years of age, respectively.

In the base case, TAVI was associated with total costs that were about DKK65,000 higher than SAVR (276,142 vs. 211,581), but TAVI patients also accumulated 0.09 additional quality-adjusted life years (QALYs; 5.39 vs. 5.30). The resulting incremental cost-effectiveness ratio (ICER) of DKK696,264/QALY (around €72,100/QALY using purchasing power parity (PPP)-adjusted conversion via U.S. dollars; 1 DKK = €0.1036) was below the willingness-to-pay threshold for cost-effective (~DKK1.1 million/QALY) and above the willingness-to-pay threshold for highly cost-effective (DKK375,489/QALY). The discounted ICER with quality-unadjusted, but still discounted, life years as effectiveness parameters was DKK409,011/life year gained. See Table 2 and Supplementary Table 6.

### Uncertainty Analyses

Sensitivity analyses revealed that the model projections were robust in terms of limited sensitivity to variation of short-term complications, except for the following parameters: relative mortality risks for TAVI and SAVR over 60 months, peri-procedural TAVI mortality, and relative risk of mortality post stroke. Figure 2 and Supplementary Figure 4 display the result of a set of key deterministic 1-way sensitivity analyses in a Tornado diagram. As per a threshold analysis, large differences in mortality risk (e.g., initial mortality RR of 0.71 for SAVR – and assumed concurrent RR for TAVI of 1.23) would push the results above the willingness-to-pay threshold for cost-effectiveness. Likewise, if the short-term absolute TAVI mortality was more than 1.7% higher than the SAVR mortality or if the relative risk of stroke was under 0.72, TAVI would not be cost-effective. Variation of all other input parameters across defined ranges did not change the decision; the largest impact was seen when assessing different TAVI procedure and

SAVR index hospitalization – in particular the length of the intensive care unit stay, as well as the length of stay on the regular ward for both procedures.

Figure 3 depicts the results of the probabilistic sensitivity analysis. About 42% of the 5,000 simulations resulted in an ICER under DKK375,489 whereas 78% of the simulations were below DKK1,126,467 (see also Supplementary Figures 5-7).

Mortality was singled out as the most important driver for cost-effectiveness. Therefore, the RR for long-term mortality (i.e., beyond the observed 60-month period) was assessed further in sensitivity and scenario analyses (see Table 2 and Supplementary Figures 8-16). As expected, the continued benefit scenario as the most optimistic scenario provided the lowest ICER (DKK334,166/QALY), while the stop-and-drop scenarios as the most conservative scenarios provided the most pessimistic results (ICERs of DKK892,000/QALY and DKK904,070/QALY for the stop-and-drop scenario with an immediate drop to life tables and with a drop to and to weighted relative risk, respectively). The fade-out scenarios, which were considered the most realistic, resulted in ICERs between those 2 extreme scenarios (DKK696,264/QALY and DKK712,188/QALY for fading out to life tables and to weighted relative risk, respectively). Importantly, the two-way sensitivity analyses (Supplementary Figures 10 and 14) indicate that the width of the “corridor” where TAVI (compared to SAVR) is highly cost-effective and not just cost-effective or is dominated by SAVR depends on which scenario is analyzed, as evidenced by the difference of the blue shapes in those Figures: while the corridor for the realistic scenario is relatively broad, the best-case scenario was more susceptible to changes to the long-term mortality,

## DISCUSSION AND LIMITATIONS

The present study is the first economic evaluation to compare first-line TAVI with SAVR in a cohort comprised primarily of patients at low surgical risk. Its results indicate that the additional resources spent on TAVI are, on average, well spent given there were no statistical difference in the underlying clinical trial (3) and the health economic profile is favorable. While TAVI is not an economically dominant or cost-saving procedure in this setting, all examined scenarios fell below the cost-effective willingness-to-pay threshold set by the World Health Organization, and in 1 scenario TAVI was highly cost-effective.

The reason for this perhaps initially counterintuitive result – that TAVI is cost-effective – is that the higher TAVI device price is partially compensated for by the lower procedural and hospitalization costs, as well as the slightly lower major adverse event rates. The only exception for this is permanent pacemaker implantation due to a high risk of conduction abnormalities such as complete heart block among TAVI patients. The incidence of permanent pacemaker placement (PPM) was 34.1% at 30 days compared with 1.6% in patients treated with SAVR<sup>3</sup>, a value that was higher than rates found in contemporary trials or registries that have studied self-expanding TAVI devices<sup>18-20</sup>. Given the costs of only around DKK15,500 (excluding hospital stay) for PPM implantation in the Danish setting<sup>16</sup>, this difference in PPM rates did not have a larger impact on the results. This might be different in other healthcare systems where PPM implantation is associated with higher costs. However, data from newer studies of self-expanding TAVI devices, including Evolut R, suggest a reduction in observed PPM rates that can be expected to improve the cost-effectiveness of TAVI<sup>21</sup>.

On the other hand, the largest drivers of the value of TAVI in the NOTION trial were associated with peri-procedural and long-term mortality risk, as well as the costs for the procedure itself and the index hospitalization.

**Mortality.** Even though 4-year results have recently been made available for the NOTION trial<sup>22</sup>, the impact of just changing the relative all-cause mortality risk of 0.05 to 0.1 (compared the general population) beyond those 4 years resulted in significant changes on the incremental QALYs; while the worst-case scenarios resulted in a difference of only 0.07 QALYs, the best-case scenario showed a benefit of around 0.19 additional QALYs for TAVI patients. It was these incremental QALY differences that drove the somewhat different ICERs, and the base care that we chose, one of the fade-out scenarios (termed “realistic” scenario), was a middle ground between the possibly too conservative stop-and-drop scenarios and the possibly too optimistic continuous-benefit scenarios.

While we varied the RRs for the long-term mortality between a large range of parameters, the 1-way analyses should be interpreted with caution. For example, the ICER of nearly DKK2.8 million/QALY gained for a SAVR RR of 0.5 would only have an effect if the TAVI RR stayed at 1.05, which is unrealistic based on the 4-year follow-up results from NOTION<sup>5</sup>. Therefore, the 2-way sensitivity analyses can be helpful, providing reassurance that even when increasing or decreasing the RR for SAVR a bit more than the one for TAVI would still result in a similar ICER range; in particular, the 2-way sensitivity analysis for the realistic scenario contained a broad cost-effectiveness “corridor.”

Most importantly, a difference in periprocedural mortality between TAVI and SAVR of 2% will render the procedure cost-ineffective. However, it is not likely that the periprocedural mortality of TAVI in other cohorts will exceed that of SAVR given that there was no statistically significant difference in the low-risk NOTION cohort<sup>3-5</sup> and the intermediate-risk SURTAVI cohort<sup>19</sup>, and superior at the initial follow-up of the CoreValve High Risk cohort<sup>18</sup>. The same holds for data observed in previous trials of balloon-expandable TAVI devices<sup>23,24</sup>.

**TAVI Procedure Costs.** While comments about future device price developments would be speculative, it is not just possible but seems highly likely that with increasing procedural experience, procedures and discharge processes will be streamlined, and other costs associated with TAVI would decrease further<sup>25,26</sup>.

**Hospitalization Costs.** The costs for the index hospitalization are considerable, DKK132,330 after SAVR, not including additional costs for various procedure-related adverse events. SAVR has been performed for decades, and discharge from both the intensive care unit and regular floors can be considered to have little possible room for efficiency gains given the maturity of the procedure, even though performance might vary between surgeons and centers<sup>27</sup>. In particular, length of stay might be slightly longer in the Nordic countries compared to the U.S. and certain other European countries. What seems sensible to point out is that after these long hospitalizations and procedure-related adverse events, old and frail patients will have a harder time recovering from “post-hospital syndrome”<sup>28</sup> – so every day less spent in the hospital, and in particular in intensive care unit settings, might be even more meaningful for the relatively old patients with severe aortic stenosis and their relatives than can be gathered in this data. On the

other hand, fast-track hospital stay and even next-day discharge has been developed for selected patients after TAVI <sup>25</sup>, and the length of stay after TAVI in the NOTION trial might have improved since the patients for this study were recruited in between 2009 and 2014.

**Surgical Risk.** Compared with cost-effectiveness analyses in high or extreme surgical risk cohorts <sup>2</sup>, the additional benefit associated with TAVI that was found in the present study is smaller. Given that the NOTION trial was a low surgical risk cohort, different results are to be expected. The relatively small effectiveness gain is in line with the clinical results from the low-<sup>3, 4, 22</sup> and intermediate-risk <sup>19</sup> cohorts and underscores that the additional expenditures for TAVI are partially compensated for by other cost-saving mechanisms, including the lower procedure costs apart from the device costs, the lower index hospitalization costs, and somewhat lower complication costs apart from PPM.

**Limitations.** Our study is subject to several limitations. First, the present analyses apply only to Denmark as different cost structures in different healthcare systems might lead to different results. Nevertheless, if potentially different cost structures would apply in a U.S. setting, then this would translate to a PPP-adjusted ICER of around \$102,000/QALY gained which, per the ACC/AHA guidelines on costs and value <sup>29</sup>, can be considered a cost-effective intervention of an intermediate value. Second, a decision-analytic model is just a representation of the reality and will never be perfect. However, modeling might help to better inform decision-makers, and it highlights areas in which uncertainty exists and possibly even quantifies this uncertainty. Third, the long-term mortality of TAVI and SAVR beyond 4 years is unclear. However, we have conducted extensive sensitivity and scenario analyses. Fourth, it is unclear whether the outcomes

of the NOTION trial are applicable to other settings, for example in patients with another distribution of access routes, surgical risks, and comorbidities. However, the NOTION trial was the first to study TAVI in a low surgical risk cohort, and insights from the Danish setting provided valuable insight for other trials and observational studies in either low- or intermediate surgical risk cohorts. Fifth, cost structures and absolute costs, in particular for the procedures itself and for the index hospitalization, tend to vary dramatically between the United States and other countries<sup>30</sup>; consequently, predictions about cost-effectiveness of self-expanding TAVI devices in other settings may prove difficult. Sixth, while some cost estimates were derived from national prices on a near-cost basis, other costs and prices were taken from a single Danish institution. However, TAVI and SAVR are performed in only four Danish centers, and costs are likely to be relatively similar between them. Finally, the results herein might be applicable only to self-expanding and not to balloon-expandable TAVI devices.

## CONCLUSION

TAVI was shown to be cost-effective in a low surgical risk cohort. However, the results are highly dependent on the long-term mortality and are applicable only to Denmark or to countries with similarly structured direct medical costs. While model-based analyses can help facilitate an appreciation for the cost and effectiveness drivers, long-term registries should be pursued to decrease the uncertainty around long-term mortality. Likewise, cost-effectiveness analyses should be conducted in other countries to prove or disapprove cost-effectiveness of TAVI in other low risk settings.

## IMPACT ON DAILY PRACTICE

TAVI might not just be an option for all patients (regardless of their surgical risk) with severe aortic stenosis in that its results are clinically not different from to SAVR, but could also be cost-effective, depending on the setting. Suitability of individual patients for a trans-catheter procedure should ultimately be decided on and the actual procedure performed by a dedicated service as both outcomes and costs might depend on the experience and procedural skill of that service.

While the cost-effectiveness of TAVI has been demonstrated for low-risk patients in Denmark, different cost structures or long-term mortality in other countries and settings, e.g. in patients at other surgical risks, in newer self-expanding or in balloon-expandable devices, might come to other conclusions regarding their health economic profiles.

## REFERENCES

1. Grover FL, Vemulapalli S, Carroll JD, Edwards FH, Mack MJ, Thourani VH, Brindis RG, Shahian DM, Ruiz CE, Jacobs JP, Hanzel G, Bavaria JE, Tuzcu EM, Peterson ED, Fitzgerald S, Kourtis M, Michaels J, Christensen B, Seward WF, Hewitt K, Holmes DR, Jr., Registry SAT. 2016 Annual Report of The Society of Thoracic Surgeons/American College of Cardiology Transcatheter Valve Therapy Registry. *J Am Coll Cardiol* 2017;**69**:1215-1230.
2. Huygens SA, Takkenberg JJ, Rutten-van Molken MP. (2017) Systematic review of model-based economic evaluations of heart valve implantations. *Eur J Health Econ*, 10.1007/s10198-017-0880-z.
3. Thyregod HG, Steinbruchel DA, Ihlemann N, Nissen H, Kjeldsen BJ, Petursson P, Chang Y, Franzen OW, Engstrom T, Clemmensen P, Hansen PB, Andersen LW, Olsen PS, Sondergaard L. Transcatheter Versus Surgical Aortic Valve Replacement in Patients With Severe Aortic Valve Stenosis: 1-Year Results From the All-Comers NOTION Randomized Clinical Trial. *J Am Coll Cardiol* 2015;**65**:2184-94.
4. Sondergaard L, Steinbruchel DA, Ihlemann N, Nissen H, Kjeldsen BJ, Petursson P, Ngo AT, Olsen NT, Chang Y, Franzen OW, Engstrom T, Clemmensen P, Olsen PS, Thyregod HG. Two-Year Outcomes in Patients With Severe Aortic Valve Stenosis Randomized to Transcatheter Versus Surgical Aortic Valve Replacement: The All-Comers Nordic Aortic Valve Intervention Randomized Clinical Trial. *Circ Cardiovasc Interv* 2016;**9**.
5. Søndergaard L. Clinical, safety and echocardiographic outcomes from the Nordic Aortic Valve Intervention (NOTION) trial: Four-year follow-up data in all-comer patients with severe aortic valve stenosis. In. *EuroPCR*. Paris, France; 2017.
6. Husereau D, Drummond M, Petrou S, Carswell C, Moher D, Greenberg D, Augustovski F, Briggs AH, Mauskopf J, Loder E. (2013) Consolidated Health Economic Evaluation Reporting Standards (CHEERS) statement. *BMJ*, 10.1136/bmj.f1049.

7. Anderson JL, Heidenreich PA, Barnett PG, Creager MA, Fonarow GC, Gibbons RJ, Halperin JL, Hlatky MA, Jacobs AK, Mark DB, Masoudi FA, Peterson ED, Shaw LJ. ACC/AHA statement on cost/value methodology in clinical practice guidelines and performance measures. *J Am Coll Cardiol* 2014;**63**:2304.
8. Drummond M, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. 9. Economic evaluation using decision-analytic modelling. In. *Methods for the Economic Evaluation of Health Care Programmes*. 4th ed. Oxford: Oxford University Press; 2015, 311-352.
9. Kappetein AP, Head SJ, Genereux P, Piazza N, van Mieghem NM, Blackstone EH, Brott TG, Cohen DJ, Cutlip DE, van Es GA, Hahn RT, Kirtane AJ, Krucoff MW, Kodali S, Mack MJ, Mehran R, Rodes-Cabau J, Vranckx P, Webb JG, Windecker S, Serruys PW, Leon MB. Updated standardized endpoint definitions for transcatheter aortic valve implantation: the Valve Academic Research Consortium-2 consensus document. *J Am Coll Cardiol* 2012;**60**:1438-54.
10. Drummond M, Sculpher MJ, Torrance GW, O'Bernie BJ, Stoddart GL. 6. Cost-utility analysis. In. *Methods for the Economic Evaluation of Health Care Programmes*. 4th ed. Oxford, UK: Oxford University Press; 2015, 137-210.
11. Bach JP, Riedel O, Pieper L, Klotsche J, Dodel R, Wittchen HU. Health-related quality of life in patients with a history of myocardial infarction and stroke. *Cerebrovasc Dis* 2011;**31**:68-76.
12. Statbank Denmark. FRDK416: Assumptions of Mortality for the Population Projection 2016 by Sex and Age. In.
13. Cleemput I, Neyt M, Thiry N, Laet CD, Leys M. *Threshold Values for Cost-Effectiveness in Health Care. KCE Reports 100C*. Brussels: Belgian Health Care Knowledge Centre; 2008.
14. Marseille E, Larson B, Kazi DS, Kahn JG, Rosen S. Thresholds for the cost-effectiveness of interventions: alternative approaches. *Bull World Health Organ* 2015;**93**:118-24.
15. International Monetary Fund, World Bank, United Nations. List of countries by GDP (nominal) per capita. In.

16. Personal Communication. Financial Department, The Heart Center, Rigshospitalet, University Hospital of Copenhagen, Denmark. In.
17. Drummond M, Sculpher MJ, Claxton K, Stoddart GL, Torrance GW. In. *Methods for the Economic Evaluation of Health Care Programmes*. 4th ed. Oxford, UK: Oxford University Press; 2015, 315-319.
18. Adams DH, Popma JJ, Reardon MJ, Yakubov SJ, Coselli JS, Deeb GM, Gleason TG, Buchbinder M, Hermiller J, Jr., Kleiman NS, Chetcuti S, Heiser J, Merhi W, Zorn G, Tadros P, Robinson N, Petrossian G, Hughes GC, Harrison JK, Conte J, Maini B, Mumtaz M, Chenoweth S, Oh JK, Investigators USCC. Transcatheter aortic-valve replacement with a self-expanding prosthesis. *N Engl J Med* 2014;**370**:1790-8.
19. Reardon MJ, Van Mieghem NM, Popma JJ, Kleiman NS, Sondergaard L, Mumtaz M, Adams DH, Deeb GM, Maini B, Gada H, Chetcuti S, Gleason T, Heiser J, Lange R, Merhi W, Oh JK, Olsen PS, Piazza N, Williams M, Windecker S, Yakubov SJ, Grube E, Makkar R, Lee JS, Conte J, Vang E, Nguyen H, Chang Y, Mugglin AS, Serruys PW, Kappetein AP, Investigators S. Surgical or Transcatheter Aortic-Valve Replacement in Intermediate-Risk Patients. *N Engl J Med* 2017;**376**:1321-1331.
20. Sorajja P, Kodali S, Reardon MJ, Szeto WY, Chetcuti SJ, Hermiller J, Jr., Chenoweth S, Adams DH, Popma JJ. Outcomes for the Commercial Use of Self-Expanding Prostheses in Transcatheter Aortic Valve Replacement: A Report From the STS/ACC TVT Registry. *JACC Cardiovasc Interv* 2017;**10**:2090-2098.
21. Grube E, Van Mieghem NM, Bleiziffer S, Modine T, Bosmans J, Manoharan G, Linke A, Scholtz W, Tchetché D, Finkelstein A, Trillo R, Fiorina C, Walton A, Malkin CJ, Oh JK, Qiao H, Windecker S. Clinical Outcomes With a Repositionable Self-Expanding Transcatheter Aortic Valve Prosthesis: The International FORWARD Study. *J Am Coll Cardiol* 2017;**70**:845-853.

22. Søndergaard L. Clinical, safety and echocardiographic outcomes from the Nordic Aortic Valve Intervention (NOTION) trial: four-year follow-up data in all-comer patients with severe aortic valve stenosis. In. *EuroPCR*. Paris, France; 2017.
23. Leon MB, Smith CR, Mack MJ, Makkar RR, Svensson LG, Kodali SK, Thourani VH, Tuzcu EM, Miller DC, Herrmann HC, Doshi D, Cohen DJ, Pichard AD, Kapadia S, Dewey T, Babaliaros V, Szeto WY, Williams MR, Kereiakes D, Zajarias A, Greason KL, Whisenant BK, Hodson RW, Moses JW, Trento A, Brown DL, Fearon WF, Pibarot P, Hahn RT, Jaber WA, Anderson WN, Alu MC, Webb JG, Investigators P. Transcatheter or Surgical Aortic-Valve Replacement in Intermediate-Risk Patients. *N Engl J Med* 2016;**374**:1609-20.
24. Leon MB, Smith CR, Mack M, Miller DC, Moses JW, Svensson LG, Tuzcu EM, Webb JG, Fontana GP, Makkar RR, Brown DL, Block PC, Guyton RA, Pichard AD, Bavaria JE, Herrmann HC, Douglas PS, Petersen JL, Akin JJ, Anderson WN, Wang D, Pocock S, Investigators PT. Transcatheter aortic-valve implantation for aortic stenosis in patients who cannot undergo surgery. *N Engl J Med* 2010;**363**:1597-607.
25. Lauck SB, Wood DA, Baumbusch J, Kwon JY, Stub D, Achtem L, Blanke P, Boone RH, Cheung A, Dvir D, Gibson JA, Lee B, Leipsic J, Moss R, Perlman G, Polderman J, Ramanathan K, Ye J, Webb JG. Vancouver Transcatheter Aortic Valve Replacement Clinical Pathway: Minimalist Approach, Standardized Care, and Discharge Criteria to Reduce Length of Stay. *Circ Cardiovasc Qual Outcomes* 2016;**9**:312-21.
26. Toggweiler S. How to reduce costs in transcatheter aortic valve implantation. In. *Open Heart*; 2014, e000203.
27. Hannan EL, Cozzens K, King SB 3rd, Walford G, Shah NR. The New York State cardiac registries: history, contributions, limitations, and lessons for future efforts to assess and publicly report healthcare outcomes. *J Am Coll Cardiol* 2012;**59**:2309-16.

28. Krumholz HM. Post-hospital syndrome—an acquired, transient condition of generalized risk. *N Engl J Med* 2013;**368**:100-2.
29. Anderson JL, Heidenreich PA, Barnett PG, Creager MA, Fonarow GC, Gibbons RJ, Halperin JL, Hlatky MA, Jacobs AK, Mark DB, Masoudi FA, Peterson ED, Shaw LJ. ACC/AHA statement on cost/value methodology in clinical practice guidelines and performance measures: a report of the American College of Cardiology/American Heart Association Task Force on Performance Measures and Task Force on Practice Guidelines. *J Am Coll Cardiol* 2014;**63**:2304-22.
30. Anderson GF, Reinhardt UE, Hussey PS, Petrosyan V. It's the prices, stupid: why the United States is so different from other countries. *Health Aff* 2003;**22**:89-105.
31. Langkilde LK, Bergholdt Asmussen M, Overgaard M. Cost-effectiveness of dabigatran etexilate for stroke prevention in non-valvular atrial fibrillation. Applying RE-LY to clinical practice in Denmark. *J Med Econ* 2012;**15**:695-703.
32. Sabale U, Ekman M, Granstrom O, Bergenheim K, McEwan P. Cost-effectiveness of dapagliflozin (Forxiga(R)) added to metformin compared with sulfonylurea added to metformin in type 2 diabetes in the Nordic countries. *Prim Care Diabetes* 2015;**9**:39-47.

## FIGURE TITLES AND LEGENDS

**Figure 1: Model Structure.** Patients start with the index procedure and associated risk of periprocedural AEs; death, stroke, MI, major or life-threatening bleeding, cardiogenic shock, acute kidney injury, major vascular complication, atrial fibrillation, permanent pacemaker implantation, prosthetic valve endocarditis, or the need for reintervention. Mortality, stroke, and MI probabilities and utilities were calculated for separately for the following states: NYHA I/II, NYHA III/IV, NYHA I/II after an MI, NYHA III/IV after an MI, NYHA I/II after a stroke, and NYHA III/IV after a stroke. All other AE consequences were included in transition probabilities and utilities for these strata, but costs occurred separately as per the calculated event probabilities.

**Figure 2: One-way Sensitivity Analyses in the Form of a Tornado Diagram.** Data categories are listed vertically instead of the standard horizontal presentation, and the categories are ordered so that the largest bar (based on the largest spread of the ICER) appears at the top of the chart, the second largest appears second from the top, and so on. A black bar represents the ICER for the high value of the varied input parameter and a white bar the low value. The base case and the willingness-to-pay threshold for “cost-effective” are marked with dashed lines.

**Figure 3: Probabilistic Sensitivity Analysis.** The left panel shows a scatter plot containing the 5,000 simulation results as sets of incremental costs (y-axis) and incremental effectiveness in QALYs (x-axis). The two willingness-to-pay (WTP) thresholds of DKK375,489 and DKK1,126,467 are marked as dashed lines. The right panel contains the corresponding willingness to pay threshold where the y-axis depicts the percentage of simulations that are cost-effective at the given WTP threshold on the x-axis

Table 1: Key Input Parameters

	<b>TAVI</b>	<b>SAVR</b>	<b>Distribution</b>	<b>Source</b>
Age (years)	79.1		Normal	3
Gender (% female)	46.2	47.4%	Beta	3
<b>Immediate AEs</b>				
Major/life-threatening bleeding	11.3%	20.9	Beta	post-hoc analysis
Cardiogenic shock	4.2%	10.4%	Beta	post-hoc analysis
Acute kidney injury	0.7%	6.7%	Beta	post-hoc analysis
Major vascular complications	5.6%	1.5%	Beta	post-hoc analysis
<b>30-day/1-year AEs</b>				
All-cause mortality	2.1% /4.9%	3.7%/7.5%	Beta	3
Stroke	1.4% /2.9%	3.0%/4.6%	Beta	3
MI	2.8% /3.5%	6.0%/6.0%	Beta	3
Atrial fibrillation	16.9%%/21.2%	57.8%/59.4%	Beta	3

PPM implantation	34.1% /38.0%	1.6% /2.4%	Beta	3
Prosthetic valve IE	0.7% /2.9%	0%/1.6%	Beta	3
Reintervention	0%/0%	0%/	n.a.	post-hoc analysis
<b>Outpatient costs</b>				
Transthoracic echocardiography	DKK 1,150		Gamma	16
Coronary angiography	DKK 7,749			16
CTA chest	DKK 2,621			16
CV specialist visit	DKK 1,362			16
<b>Procedure costs</b>				
Procedure time (minutes)	90	177		16
Physician time	DKK 815	DKK 1,597		skin-to-skin time; wages <sup>16</sup>
Nursing time	DKK 415	DKK 813		skin-to-skin time; wages <sup>16</sup>
Tech time		DKK 813		skin-to-skin time; wages <sup>16</sup>
Valve costs	DKK 125,000	DKK 14,906		16
Heart-lung machine	n.a.	DKK 9,600		16
Concomitant CABG		DKK 25,000		16

*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*


Other materials	DKK 9,600	DKK 6,700		16
Blood products	DKK 227	DKK 866		16
Total procedure costs	DKK 136,057	DKK 31,189	Gamma	16
<b>AE costs</b>				
Major or life-threatening bleeding	DKK 4,752		Gamma	national price, <sup>16</sup>
Cardiogenic shock	DKK 29,638		Gamma	national price, <sup>16</sup>
Acute kidney injury	DKK 11,342		Gamma	national price, <sup>16</sup>
Major vascular complications	DKK 12,171		Gamma	national price, <sup>16</sup>
Stroke: first month	DKK 34,711		Gamma	31, 32
Stroke: remaining months	DKK 3,192		Gamma	31, 32
MI: first month	DKK 19,085		Gamma	31, 32
MI: remaining months	DKK 136		Gamma	31, 32
Atrial fibrillation	DKK 6,510		Gamma	national price, <sup>16</sup>
PPM implantation	DKK 15,500		Gamma	16
Prosthetic valve IE	DKK 80,586		Gamma	16

*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*


Atrial fibrillation	DKK 11,972		Gamma	national price, <sup>16</sup>
PPM implantation	DKK 15,000		Gamma	<sup>16</sup>
prosthetic valve IE	DKK 80,586		Gamma	national price, <sup>16</sup>
<b>LOS and related costs</b>				
ICU LOS (days)	1.2	2.6	Gamma	<sup>16</sup>
Regular ward LOS (days)	7.7	10.3	Gamma	<sup>16</sup>
Costs per ICU day	DKK 22,915		Gamma	<sup>16</sup>
Costs per regular ward day	DKK 5,462		Gamma	<sup>16</sup>
Total index hospitalization	DKK 76,109	DKK 132,330	Gamma	

**Table 2: Results of the Base Case and Other Long-term Mortality Scenarios**


	Costs (DKK)	Incremental Costs (DKK)	QALYs	Incremental QALYs	ICER (DKK/QALY)
<b>Realistic Scenario I (base case – fade out over 24 months to life tables [RR=1])</b>					
SAVR	211,581		5.30		
TAVI	276,142	64,561	5.39	0.09	696,264
<b>Realistic Scenario II (fade out over 24 months to weighted RR [1.10] – then continues lifelong)</b>					
SAVR	210,826		5.19		
TAVI	275,329	64,503	5.28	0.09	712,188
<b>Best-case Scenario I (group-specific RRs [1.05/1.15] continue lifelong)</b>					
SAVR	210,484		5.14		
TAVI	275,722	64,238	5.33	0.19	334,166
<b>Worst-case Scenario I (immediate drop to life tables [RR=1.0])</b>					
SAVR	211,881		5.33		
TAVI	276,238	64,357	5.40	0.07	892,000
<b>Worst-case Scenario II (immediate drop to weighted RR [1.10] – but continues lifelong)</b>					
SAVR	210,920		5.20		
TAVI	275,239	64,319	5.27	0.07	904,070




*Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal*



**Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal**



**Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal**



**Disclaimer : As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal**

## SUPPLEMENTARY MATERIALS

for

Cost-Effectiveness of Transcatheter versus Surgical Aortic Valve Implantation in Patients at Lower Surgical Risk

Results from the NOTION Trial

Benjamin P. Geisler, M.D.<sup>1,2</sup>, Troels H. Jørgensen<sup>3</sup>, Hans Gustav H. Thyregod, M.D., Ph.D.<sup>3</sup>, Jan B. Pietzsch, PhD.<sup>1</sup>, and Lars Søndergaard, M.D., Ph.D.<sup>3</sup>

From <sup>1</sup>Wing Tech Inc., Menlo Park, CA, USA; <sup>2</sup>Massachusetts General Hospital/Harvard Medical School, Boston, MA, USA; <sup>3</sup>Rigshospitalet, Copenhagen, Denmark

Correspondence to: Jan B. Pietzsch, Ph.D., Wing Tech Inc., 101 Jefferson Drive, Menlo Park, CA 94025, USA; [jpietzsch@wing-tech.com](mailto:jpietzsch@wing-tech.com), fax +1 (301) 983-2558, phone +1 (650) 943-3000

Classifications: TAVI; Death; No specific risk; Miscellaneous.

Copyright EuroIntervention

## Methods

### Model Structure and Decision-Analytic Framework

The decision-analytic model was constructed in TreeAge 2019 R1.1 (TreeAge Software, Inc., Waltham, MA) to forecast long-term societal costs and effectiveness and to compute an incremental cost-effectiveness ratio (ICER). The model was a combination of a decision tree and a state transition or Markov model where the patients' disease progression through defined, mutually exclusive health states is tracked at monthly intervals<sup>8</sup>. The model was run with half-cycle correction until the entire cohort was deceased (lifetime horizon).

All patients started off with the index procedure, TAVI or SAVR, where they could experience the following Valve Academic Research Consortium-2-defined periprocedural events<sup>9</sup>: all-cause death, fatal and nonfatal stroke, fatal and nonfatal MI, major or life-threatening bleeding, cardiogenic shock, stage II-III acute kidney injury, major vascular complication, atrial fibrillation, permanent pacemaker implantation (PPM), prosthetic valve infective endocarditis, or the need for reintervention. These events were accounted for in terms of costs and mortality. Utilities, derived from trial-collected SF-6D data, were estimated to adjust life expectancy for health-related quality of life<sup>10</sup>.

For modeling purposes, the study population was stratified into the following subgroups: NYHA I/II or III/IV without stroke or MI, with stroke, or with MI, respectively, based on the assumption that these events would impact both costs and health-related quality of life<sup>11</sup>. If the patients did not die during the procedure, we calculated their monthly probability of experiencing NYHA I/II vs. III/IV symptoms and either an MI or a stroke (see below and Figure 1 in the main manuscript for details). In addition, we also accounted for the following 1-year adverse events in terms of costs: atrial fibrillation, PPM, and infective endocarditis. For the long-term mortality beyond the observed trial period, we defined as a reference the Danish life tables<sup>12</sup>. This means that a relative risk (RR) of 1.0 corresponds to the mortality of the general population. On top of this background mortality derived from the life tables, various scenarios analyzing

different combinations of RRs for TAVI and SAVR were modeled (see below for details). A simplified representation of the model structure is shown in Figure 1 in the main manuscript.

In the absence of other clear recommendations for Denmark<sup>13</sup>, the main decision criterion to evaluate cost-effectiveness was whether the ICER was below the willingness-to-pay thresholds established by the World Health Organization<sup>14</sup>. These thresholds stipulate that an intervention is “cost-effective” if the ICER is less than 3 times the gross domestic product per capita of the respective country and “highly cost-effective” if below the per-capita gross domestic product. All costs in this analysis are expressed in 2016 Danish kroner (DKK) and hence used the nominal 2016 gross domestic product per capita of DKK375,489<sup>15</sup>. The discount rate was 3% for costs and effectiveness.

### Patient Characteristics and Other Input Parameters

Patient characteristics were based on the only available trial of TAVR vs SAVR in low-risk patients, NOTION 3; at baseline, there were 280 patients, mean age at the time of the procedure was 79.1 ±4.8 years, 46.8% of the cohort were female, and their mean Society of Thoracic Surgeons Predicted Risk of Mortality and logistic EuroSCORE II were 3.0% and 2.0%, respectively. 96.5% of patients had transfemoral access. Pre-operative test and consults included transthoracic echocardiography, coronary angiography, and visits to a cardiologist and/or a cardiothoracic surgeon (the latter for SAVR patients only). Computed tomography for sizing of the aortic valve annulus was not routinely performed. For TAVI, only the CoreValve self-expandable bioprosthesis was used (Medtronic, Minneapolis, MN), whereas surgical patients received any commercially available bioprosthetic aortic valve at the discretion of the surgeon. In 2.8% of patients, two valve kits with the corresponding costs were assumed. The 30-day mortality in NOTION was 2.1% vs. 3.7% for TAVI vs. SAVR, respectively; 1.4% and 3.0% experienced a stroke, and 2.8% and 6.0% an MI. At 1 year, 92.4% vs. 96.4% of the patients, respectively, had NYHA I/II class symptoms. A TAVI device cost of DKK125,000 was assumed<sup>16</sup>, and for SAVR a bioprosthetic and not a mechanical valve at DKK14,906 without the need for systemic anticoagulation for

all patients given their age; all device costs were subsumed in the procedure costs. In the NOTION trial, patients had a mean intensive care unit length of stay of 1.3 vs. 2.6 days after TAVI and SAVR and remained on a general ward for 8.9 and 12.9 days, respectively<sup>3</sup>. Amid these lengths of stay, none of the patients was assumed to have been transferred from the hospital to a free-standing physical rehabilitation center at the time of discharge.

Monthly probabilities of experiencing all-cause death, stroke, or MI in the remaining patients at risk were computed from the patient-level NOTION 5-year data set<sup>4,5</sup>. All statistical analyses were conducted in SAS 9.4 (SAS Institute, Cary, NC). Resulting key input parameters for the model in the Appendix Table 1.

## Model Calibration and Uncertainty Analyses

To achieve concordance with the actual trial results, the present model was calibrated with regard to the following endpoints: all-cause mortality, stroke, or MI. See below for details on the calibration.

To evaluate the effects of uncertainty in input parameters, we conducted the following analyses: For all input parameters, 1-way sensitivity analyses were performed by changing the input parameter by at least  $\pm 30\%$ . The input parameters to which the model was most sensitive (as judged by the spread of net monetary benefit) were summarized in Tornado diagrams. We also performed a probabilistic sensitivity analysis with 5,000 random draws in a second-order Monte Carlo simulation (see further down below).

Extensive sensitivity analyses were performed regarding the long-term mortality as this was *a posteriori* judged to be one of the most important parameters for a low-risk cohort (see Results in the main manuscript and below). The RRs were analyzed in separate 1-way sensitivity analyses and in a 2-way analysis. Several scenarios were created in accordance with established standards on how to extrapolate along-the-trial results<sup>17</sup>: First, 2 “stop-and-drop” scenarios (most conservative scenarios), one involving dropping the RRs to 1.0 (i.e., applying the mortality from the life tables of the general population to both TAVI and SAVR arms) and one dropping it to the weighted mortality observed in the second 2-year

interval of the trial (combined RR: 1.10, applied to both arms). Second, the most optimistic scenario was a “continued-benefit” scenario where the mortality benefit of TAVI over SAVR observed in the last 2 years of the trial continued for the remainder of the patients’ lifetime. Third, for the base case, we used a “fade-out” scenario, where the relative risks of TAVI and SAVR both were subject to a linear fading out over 24 months toward a RR of 1.0, i.e., the mortality of the 2 groups was approaching the life tables of the general population over this time. A second fade-out scenario examined the effect of a fade-out from the observed RRs (RR for TAVI: 1.05; RR for SAVR: 1.15) toward the weighted combined RR of 1.10 as compared to general population lifetables. In addition, selected threshold analyses were conducted.

### Complete List of Input Parameters

In Supplementary Table 1, we list all input parameters used in the decision-analytic model. The parameters are distributed into the following categories (as they are in all sections below):

- Patient characteristics
- Probability of immediate adverse events
- Probability of 30-day adverse events
- Probability of 1-year adverse events
- Preoperative costs
- Procedure costs
- Costs per immediate adverse event
- Length of stay and hospitalization-related costs
- Costs per 30-day adverse events
- Costs of follow-up care
- Costs per 1-year adverse events

Some parameters are different for the TAVI and the SAVR arm and are listed in the respected column.

The values in the brackets correspond to the ranges for the deterministic sensitivity analyses. The second row lists the distribution assumed in the probabilistic sensitivity analysis with the corresponding parameters.

## Trial-based Mortality, Stroke, and MI Incidence, and Model Calibration

This section shows a set of figures with an overlay of the calibrated model output (all-cause mortality, stroke, and myocardial infarction) over the actual trial outcome with the corresponding patients at risk in the TAVI and SAVR groups, respectively. The trial outcomes were taken from the publications and presentations describing the outcomes of the NOTION study<sup>3-5</sup> as well as the five-year outcomes which are currently in print. The model output was generated via Markov traces for each treatment strategy separately.

### All-cause Mortality

Supplementary Figure 1 depicts all-cause mortality. The actual TAVI trial outcome is represented the orange line and the TAVI model output by the red circles. The actual SAVR trial outcome is represented by the turquoise line and the SAVR model output by the green circles.

The reported trial outcome and calibrated model output at months one, twelve, 24, 48, and 60 is summarized again in the **Error! Reference source not found.**

Of note, these model results match the actual trial because we manually calibrated the transition probabilities to the actual trial results.

### Stroke

Supplementary Figure 2 depicts the incidence of stroke: The actual TAVI trial outcome is represented by the orange line and the calibrated model output by the red circles. The actual SAVR trial outcome is represented by the turquoise line, and the green circles represent the model output.

The reported trial outcome and calibrated model output in terms of stroke incidence at months 1, 12, 24, 48, and 60 is summarized again in **Error! Reference source not found.**

Of note, these model results match the actual trial because we manually calibrated the transition probabilities to the actual trial results.

## Myocardial Infarction

Supplementary Figure 3 depicts the incidence of myocardial infarction: The actual TAVI trial outcome is represented by the orange line and the calibrated model output by the red circles. The actual SAVR trial outcome is represented by the turquoise line, and the green circles represent the model output.

The reported trial outcome and calibrated model output in terms of stroke incidence at months one, twelve, 24, 48, and 60 is summarized again in **Error! Reference source not found.**

Of note, these model results match the actual trial because we manually calibrated the transition probabilities to the actual trial results.

## Utilities

Supplementary Table 2 to Supplementary Table 5 summarize the utilities used in the model, based on scored SF-6D questionnaires collected in NOTION.

## Probabilistic Sensitivity Analysis

A probabilistic sensitivity analysis was performed to quantify the overall uncertainty. We used 54 distributions for the input parameters (see table 1 in the main manuscript). In a second-order Monte Carlo simulation, 5,000 random draws from these distributions were performed, and the model was run to estimate a cost/effectiveness data point for TAVI and SAVR, which are depicted on the two subsequent incremental cost-effectiveness (ICE scatter plots). The ellipse on each represents the area where 95% of the simulations are located.

Supplementary Figure 5 demonstrates the results of the probabilistic sensitivity analysis with a willingness-to-pay threshold of DKK 375,489/QALY which is equivalent to “highly cost-effective”.

Supplementary Figure 6 demonstrates the results of the probabilistic sensitivity analysis with a willingness-to-pay threshold of DKK 1,126,467/QALY which is equivalent to “cost-effective”.

Supplementary Figure 7 is a cost-effectiveness acceptability curve. About 32% of the 5,000 simulations resulted in an ICER under DKK 375,489 (corresponding to the WHO definition of “highly cost-effective”) whereas 56% of the simulations were below DKK 1,126,467 (“cost-effective”).

### **Tornado Diagram**

Supplementary Figure 4 shows a slightly more comprehensive Tornado diagram compared to the figure in the manuscript. However, as can be seen and as discussed in the manuscript, the main drivers of the ICER are (in descending order)

1. Relative mortality risk for SAVR beyond the 48-month trial observation period (and, to a lesser degree, for the relative risk for TAVI and for the five scenarios [as discussed in extensive detail below])
2. Periprocedural mortality during TAVI (and, to a lesser degree, for corresponding periprocedural mortality of SAVR)
3. Procedure costs for TAVI (and, to a lesser degree, for SAVR procedural costs)
4. Length of stay for SAVR and TAVI index hospitalization, in particular for the ICU component of the SAVR and the regular ward component of the TAVI index hospitalizations, respectively – or costs *in toto* for the entire hospitalization or per ICU day
5. Discount rates for effectiveness

Again, we discuss these factors in the manuscript. However, a more extensive discussion on the relative mortality risks and the scenarios created from varying these in different ways can be found below.

### **Sensitivity Analyses on Long-term Mortality**

As discussed in the main manuscript and in line with the health economic literature<sup>8</sup>, we analyzed several scenarios for the long-term mortality beyond the observed 48 months:

1. Base case: Fade-out Scenario I (fade out over 12 months to life tables [RR=1])

2. Fade-out Scenario II (fade out over 12 months to weighted RR [1.10] – then continues lifelong)
3. Continuous Benefit Scenario I (group-specific RRs [1.05/1.15] continue lifelong)
4. Stop-and-Drop Scenario I (immediate drop to life tables [RR=1.0])
5. Stop-and-Drop Scenario II (immediate drop to weighted RR [1.10] – but continues lifelong)

What might be perceived as slightly different from the standard approach<sup>8</sup> is that we have two fade-out and two stop-and-drop scenarios. The reason for this is that we consider the relative risks to fade out – or drop immediately – to different values: one, to the background mortality of the general population (i.e., a relative risk of 1.0), and two, to a weighted RR of 1.1, which corresponds to a slightly worse mortality than the general population. This relative risk was derived from the weighted relative risks from the second 2 years of the NOTION trial<sup>3</sup>: To explain, we obtained the “weighted” relative risk of 1.1 from the second 2 years of the trial (months 24 to 47), which was 12.0% for TAVI and 13.2% for SAVR. This compared to a gender-adjusted mortality over the corresponding 2 years from the Danish life tables (ages 81 and 82, based on the mean age of around 79 at baseline) of 11.48%. Hence, the relative risks of TAVI and SAVR are 1.05 and 1.15. Weighing these two with the patients at risk at month 24 (n=75 and 72, respectively) results in a relative risk of 1.15.

### Fade-Out Scenarios – Base Case

In the first fade-out scenario, we let the relative mortality risk of TAVI and SAVR that was observed in the second 2 years of the trial “fade out” of 12 months to the background mortality (Danish life tables).

In the second fade-out scenario, we let the relative mortality risk of TAVI and SAVR that was observed in the second 2 years of the trial “fade out” of 24 months – as explained above.

Supplementary Figure 8 contains an illustration of the fade-out of the relative mortality risk (reference: life tables) over the first year after the observed trial period.

The sensitivity analyses regarding the long-mortality RR for the fade-out scenario are displayed in Supplementary Figure 9 and Supplementary Figure 10. Of note, we only analyzed these for the first fade-

out scenario, where the relative risk fades out to the background mortality as this had been chosen as the base case.

- Supplementary Figure 9 contains two one-way sensitivity analyses with the outcome undiscounted (and quality-*unadjusted*) life years. The first curve, in orange, analyzes the relative mortality risk (with life tables of the general population as a reference) of TAVI after the trial observation period of 48 months, the other curve, in green, the respective relative mortality risk of SAVR. Each relative risk corresponds to the x axis, and the y axis represents the outcome undiscounted life years. The base case is marked with a red circle (TAVI) and a yellow diamond (SAVR).
- Supplementary Figure 10 contains a two-way sensitivity analyses of the same relative risks, but instead of each relative risk being on the same axis (as in Supplementary Figure 9), they are represented by the x and y axes, and the color or the area corresponds to the category that the result for that respective pair of x and y values belongs to (red for TAVI being highly cost-effective, blue for TAVI being cost-effective [but not *highly* cost-effective], and purple for not cost-effective, i.e., TAVI is economically dominated by SAVR). It is important to note that when varying one RR in these sensitivity analyses, the other is fixed. For example, when assuming a SAVR RR of 0.9 in Supplementary Figure 13, the TAVI RR is kept at 1.05. Hence, these graphs represent extreme cases.

### Continuous Benefit Scenario

In the continuous benefit scenario, the estimated relative mortality risks (RRs, see above for definitions) continue lifelong. Since we observed a slightly more favorable relative risk for TAVI compared to SAVR (1.05 vs 1.1), we call this scenario “continuous benefit,” although the relative risk over 1.0 (reference: life tables) indicates that the mortality is slightly worse compared to the general population. Given that the benefit of TAVI over SAVR was observed in the last 2 years of the trial, we chose to analyze this scenario even though it might be the least conservative assumption – and is hence not the base case.

Below are several graphs that analyze the RRs as one-way sensitivity analyses between the extremes of 0.5 and 1.3 (the extremes observed in individual 12-month periods in the trial):

- Analogous to Supplementary Figure 11 (please see for an explanation above) contains two one-way sensitivity analyses with the outcome undiscounted (and quality-*unadjusted*) life years. What can be gathered immediately is that the curves are much less flat than in the fade-out scenario I.

- Supplementary Figure 12 and Supplementary Figure 13 show separate one-way sensitivity analyses on the RRs for the outcome discounted incremental cost-effectiveness ratio (ICER). Again, it is important to note that when varying one RR in these sensitivity analyses, the other is fixed. For example, when assuming a SAVR RR of 0.9 in Supplementary Figure 13, the TAVI RR is kept at 1.05. Hence, these graphs represent extreme cases.

Supplementary Figure 14 contains a two-way sensitivity analyses of the same relative risks, but instead of each relative risk being on the same axis (as in Supplementary Figure 11), they are represented by the x and y axes, and the color or the area corresponds to the category that the result for that respective pair of x and y values belongs to (red for TAVI being highly cost-effective, blue for TAVI being cost-effective [but not *highly* cost-effective], and purple for not cost-effective, i.e., TAVI is economically dominated by SAVR).

### Summary and Comparison of the Base Case Scenarios

Reproduced in Supplementary Table 6 are the results for the various scenarios. Below, please find the description to several charts to compare the different scenarios graphically.

In Supplementary Figure 15, the absolute QALYs for TAVI and SAVR are graphed by scenario. As can be seen, the stop-and-drop I scenario (to life tables) had the highest TAVI and SAVR QALY gains, followed by the fade-out I scenario (to life tables). This can of course be explained that the relative risk drops the soonest to the background mortality of the general population in the stop-and-drop scenario, but the fade-out scenario I achieves this relative risk after just 60 months. However, and more importantly, the continuous benefit scenario offers the greatest *incremental* benefit.

As can be seen in Supplementary Figure 16, the continuous benefit, which was also *a priori* believed to be the least conservative scenario, has the highest QALY gain and the lowest ICER. Noteworthy are three points:

1. The ICER largely depends on the incremental QALY difference and not on the incremental costs (these are largely the same for the different scenarios). This is likely due to the effect that 4 months out from the procedure, the most important costs are the

outpatient follow-up costs, which are similar between the strategies if survival is not much different.

2. The difference between the continuous benefit and the fade-out scenarios (which lie in the “middle”) is much greater than between the fade-out scenarios and the stop-and drop scenarios.
3. The difference between the two fade-out scenarios and the two stop-and-drop scenarios is relatively small. This is despite the higher/lower QALYs that the two strategies achieve (as can be seen in Supplementary Figure 15) but *because* of similar incremental QALY gains (as can be gathered from Supplementary Figure 16).

### Supplementary Figure Legends

Supplementary Figure 1: 60-month Mortality Data with Calibrated Model Output.....	53
Supplementary Figure 2: 60-month Stroke Incidence with Calibrated Model Output.....	53
Supplementary Figure 3: 60-Month Myocardial Infarction Incidence with Calibrated Model Output .....	54
Supplementary Figure 4: Tornado Diagram.....	55
Supplementary Figure 5: Incremental Cost-Effectiveness (ICE) Scatter PLOT with a Willingness-to-Pay Threshold of DKK 375,489 .....	56
Supplementary Figure 6: Incremental Cost-Effectiveness (ICE) Scatter PLOT with a Willingness-to-Pay Threshold of DKK 1,126,467 .....	56
Supplementary Figure 7: Cost-effectiveness Acceptability Curve.....	57
Supplementary Figure 8: Illustration of the Relative Mortality Risk Fading Out Over Time .....	57
Supplementary Figure 9: One-way Sensitivity Analyses of Relative Mortality Risk after TAVI and SAVR in the Fade-Out I Scenario (Base Case, Outcome: Undiscounted Life Years) .....	58
Supplementary Figure 10: Two-way Sensitivity Analysis on Relative Mortality Risk after SAVR over Relative Mortality Risk after TAVI (Fade-out Scenario).....	58
Supplementary Figure 11: One-way Sensitivity Analyses of Relative Mortality Risk after TAVI and SAVR in the Continuous Benefit Scenario (Outcome: Undiscounted Life Years).....	59
Supplementary Figure 12: One-way Sensitivity Analysis over Relative Mortality Risk after TAVI (Continuous Benefit Scenario).....	59
Supplementary Figure 13: One-way Sensitivity Analysis Over Relative Mortality Risk after SAVR (Continuous Benefit Scenario).....	60
Supplementary Figure 14: Two-way Sensitivity Analysis on Relative Mortality Risk after SAVR over Relative Mortality Risk after TAVI (Continuous Benefit Scenario).....	60
Supplementary Figure 15: Bar Chart Comparing the QALYs Incurred by TAVI and SAVR.....	61
Supplementary Figure 16: Combined Bar and Line Chart Comparing the Incremental QALYs (blue bars) and corresponding ICERs (orange line) .....	61

## Supplementary Tables

Supplementary Table 1: Complete List of Input Parameters

	TAVI	SAVR	Source
<b>Patient characteristics</b>			
Age (years)	79.1 (76; 82) Normal; 79.1; 4.9		3
Gender (% female)	46.8% (0%; 100%) Beta; 11.18; 12.71		3
<b>Immediate AEs</b>			
Major/life-threatening bleeding	11.3% (7.9; 14.7%) Beta; 1.02; 8.00	20.9% (14.6%; 27.2%) Beta; 3.25; 12.29	post-hoc analysis
Cardiogenic shock	4.2% (2.9; 5.5%) Beta; 0.13; 2.90	10.4% (7.3%; 13.5%) Beta; 3.25; 12.29	post-hoc analysis
Acute kidney injury	0.7% (0.5%; 0.9%) Beta; $6.03 \cdot 10^{-4}$ ; $8.55 \cdot 10^{-2}$	6.7% (4.7%; 8.7%) Beta; 0.59; 8.18	post-hoc analysis
Major vascular complications	5.6% (3.9%; 7.3%) Beta; 0.24; 4.05	1.5% (1.1%; 2.0%) Beta; $7.16 \cdot 10^{-3}$ ; 0.47	post-hoc analysis
<b>30-day AEs</b>			
All-cause mortality	2.1% Beta; $2.22 \cdot 10^{-2}$ ; 1.03	3.7% Beta; $9.48 \cdot 10^{-2}$ ; 2.47	3

Stroke	1.4% Beta; $5.33 \cdot 10^{-3}$ ; 0.38	3.0% Beta; $5.73 \cdot 10^{-2}$ ; 1.85	<sup>3</sup>
Myocardial infarction	2.8% Beta $4.82 \cdot 10^{-2}$ ; 1.67	6.0% Beta; 0.28; 4.36	<sup>3</sup>
Atrial fibrillation	16.9% (11.8%; 22.0%) Beta; 0.42; 2.09	57.8% (40.5%; 75.1%) Beta; 2.95; 2.15	<sup>3</sup>
PPM implantation	34.1% (24.0%; 44.3%) Beta 1.57; 3.04	1.6% (1.1%; 2.1%) Beta; $9.19 \cdot 10^{-3}$ ; 0.57	<sup>3</sup>
Prosthetic valve IE	0% (0.5%; 0.9%) Beta; $6.03 \cdot 10^{-4}$ ; $8.55 \cdot 10^{-2}$	0% (0%; 0.9%) Beta; $1.55 \cdot 10^{-2}$ ; 155.2	<sup>3</sup>
Reintervention	0% (0%; 5%)	0% (0%; 5%)	post-hoc analysis
<b>1-year AEs</b>			
All-cause mortality	4.9%	7.5%	<sup>3</sup>
Stroke	2.9%	4.6%	<sup>3</sup>
Myocardial infarction	3.5%	6.0%	<sup>3</sup>
Atrial fibrillation	21.2% (14.8%; 27.6%) Beta; 3.33; 12.38	59.4% (41.6%; 77.2%) Beta 13.73; 9.39	<sup>3</sup>
PPM implantation	3.9% (2.7%; 5.1%)* +34.1% (at month 1) Beta; 8.57; 13.99	0.8% (0.6%; 1.0%)* +1.6% (at month 1) Beta; $3.22 \cdot 10^{-2}$ ; 1.31	<sup>3</sup>
Prosthetic valve IE	2.9% (2.0%; 3.7%) Beta; 0.10; 3.30	1.6% (1.1%; 2.1%) Beta; $2.34 \cdot 10^{-2}$ ; 1.44	<sup>3</sup>

<b>Pre-operative Costs</b>			
Transthoracic echo	DKK 1,150		16
Coronary angiography	DKK 7,749		16
CTA chest		DKK 2,621	16
Cardiologist visit	DKK 1,362		16
Cardiothoracic surgeon visit		DKK 1,362	16
Total preoperative costs	DKK 14,244 (9,971; 18,517) Gamma; 32.46; 2.28 · 10 <sup>-3</sup>	DKK 11,623 (8,136; 15,110) Gamma; 21.62; 1.86 · 10 <sup>-3</sup>	
<b>Procedure Costs</b>			
Procedure time	90 minutes	177 minutes	16
Physician time	DKK 815	DKK 1,597	skin-to-skin time; wages <sup>16</sup>
Nursing time	DKK 415	DKK 813	skin-to-skin time; wages <sup>16</sup>
Tech time		DKK 813	skin-to-skin time; wages <sup>16</sup>
Valve costs	DKK 125,000	DKK 14,500	16
Heart-lung machine	n.a.	DKK 9,600	16
Concomitant CABG		DKK 25,000**	16
Other materials	DKK 9,600	DKK 6,700	16
PRBC transfusions	DKK 172	DKK 455	16

**Disclaimer :** As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal

FFP transfusions	DKK 18	DKK 63	16
Platelet transfusions	DKK 37	DKK 348	16
Total procedure costs	DKK 136,057*** (95,240; 176,874) Gamma; 185.12; $1.36 \cdot 10^{-3}$	DKK 31,189**** (21,832; 40546); Gamma; 155.64; $4.99 \cdot 10^{-3}$	16
<b>AE Costs During Index Hospitalization</b>			
Stroke	see below		31, 32
Myocardial infarct	see below		31, 32
Major or life-threatening bleeding	DKK 4,752 (3,326; 6,178) Gamma; 3.61; $7.60 \cdot 10^{-4}$		16
Cardiogenic shock	DKK 29,638 (20,747; 38,529) Gamma; 0.98; $3.29 \cdot 10^{-5}$		16
Acute kidney injury	DKK 11,342 (7,939; 14,745) Gamma; 0.57; $5.04 \cdot 10^{-5}$		16
Major vascular complications	DKK 12,171 (8,520; 15,822) Gamma; 5.93; $4.87 \cdot 10^{-4}$		16
Stroke: fist month	DKK 34,711 (24,298; 45,124) Gamma; 5.35; $1.54 \cdot 10^{-4}$		31, 32
Stroke: monthly costs for remaining lifetime	DKK 3,192 (2,234; 4,150) Gamma 40.76; $1.28 \cdot 10^{-4}$		31, 32
MI: fist month	DKK 64,200 (44,940; 83,460) Gamma; 3.64, $1.91 \cdot 10^{-4}$		31, 32

MI: monthly costs for remaining lifetime	DKK 136 (95; 176) Gamma; 29.59; 0.22	31, 32
Atrial fibrillation	DKK 6,510 (4,577; 8,463) plus 1.5 days LOS Gamma; 18.84; $2.89 \cdot 10^{-3}$	
PPM implantation	DKK 15,500 (21,830; 40,541) Gamma; 2.40; $1.55 \cdot 10^{-4}$	Rigshospitalet 2016
<b>LOS and related Costs</b>		
ICU LOS	1.2 days (0; 3) Gamma 6.76; 5.20	2.6 days (1; 5) 6.76; 2.60 Rigshospitalet 2016
Regular ward LOS	7.7 days (0; 10.8) Gamma 6.59; 0.86	10.3 days (3.4; 15.4) Gamma; 16.97; 1.65 Rigshospitale, 2016
Costs per ICU day	DKK 22,915 (16,041; 29,790) Gamma; 84.02; $3.67 \cdot 10^{-3}$	Rigshospitalet 2016
Costs per regular ward day	DKK 5,462 (3,823; 7,101) Gamma; 13.26; $2.43 \cdot 10^{-3}$	Rigshospitalet 2016
<b>AE Costs After Index Hospitalization</b>		
Major or life-threatening bleeding	DKK 15,676 (10973; 20379) Gamma; 39.32; $2.51 \cdot 10^{-3}$	16
Major vascular complications	DKK 12,171 (8,520; 15,822) Gamma; 31.47; $5.61 \cdot 10^{-4}$	16
Stroke: fist month	DKK 34,711 (24,298; 45,124) Gamma; 5.35; $1.54 \cdot 10^{-4}$	31, 32

Stroke: monthly costs for remaining lifetime	DKK 3,192 (2,234; 4,150) Gamma 40.76; $1.28 \cdot 10^{-4}$	31, 32
MI: fist month	DKK 64,200 (44,940; 83,460) Gamma; 3.64, $1.91 \cdot 10^{-4}$	31, 32
MI: monthly costs for remaining lifetime	DKK 136 (95; 176) Gamma; 29.59; 0.22	31, 32
Atrial fibrillation	DKK 6,510 (4,577; 8,463) Gamma; 18.84; $2.89 \cdot 10^{-3}$	
PPM implantation	DKK 15,500 plus 1.5 days LOS on a regular ward (21,830; 40,541) Gamma; 2.40; $1.55 \cdot 10^{-4}$	Rigshospitalet 2016
Prosthetic valve IE	DKK 80,586 (56,410; 104,762) Gamma; 10.39; $1.29 \cdot 10^{-4}$	
Reintervention	DKK 11,057 (7,740; 14,374)	DKK 31,189 (21,832; 40,546) calculation
<b>Follow-up Care Costs</b>		
Cardiologist visit	DKK 1,362	
Transthoracic echo	DKK 1,150	
Total monthly cost	DKK 209 (147; 272) Gamma; 17.47; 0.08	
<b>Long-term Mortality</b>		
Relative risk, compared to baseline mortality per life tables	1.05 Log normal; $3.92 \cdot 10^{-2}$ ; 0.14	1.15 Log normal; 0.12; 0.19

Listed are base case values and ranges for sensitivity analyses. AEs = adverse events; angio = angiography; CTA = computed tomography angiography; echo = echocardiography; IE = infective endocarditis; SAVR = surgical aortic valve replacement; TAVI = transcatheter aortic valve implantation.

\* These probabilities are the differences between the absolute 1-month and 1-year probabilities.

\*\* Event probability was 0.7%.

\*\*\* Includes costs for CoreValve device.

\*\*\*\* Cost does not include concomitant CABG costs (DKK 187 per case).

Copyright EuroIntervention

Supplementary Table 2: SF-6D and NYHA at 1-Month Visit

SF-6D	NYHA 1-2				NYHA 3-4			
	Major/Minor Stroke Before 1 Month		No Stroke Before 1 Month		Major/Minor Stroke Before 1 Month		No Stroke Before 1 Month	
	TAVR (N=1)	SAVR (N=2)	TAVR (N=121)	SAVR (N=106)	TAVR (N=0)	SAVR (N=0)	TAVR (N=10)	SAVR (N=4)
n	1	2	102	78	0	0	8	3
Mean ± SD	0.60 ±	0.44 ± 0.17	0.70 ± 0.12	0.66 ± 0.10	NA	NA	0.57 ± 0.11	0.56 ± 0.08
Median	0.60	0.44	0.70	0.66	NA	NA	0.59	0.60
Min, Max	0.6, 0.6	0.3, 0.6	0.4, 1.0	0.4, 0.9	NA	NA	0.4, 0.7	0.5, 0.6
Q1, Q3	0.60, 0.60	0.32, 0.57	0.62, 0.78	0.59, 0.74	NA	NA	0.49, 0.63	0.47, 0.61

Supplementary Table 3: SF-6D and NYHA at 6-Month Visit

SF-6D	NYHA 1-2				NYHA 3-4			
	Major/Minor Stroke Before 6 Month		No Stroke Before 6 Month		Major/Minor Stroke Before 6 Month		No Stroke Before 6 Month	
	TAVR (N=3)	SAVR (N=5)	TAVR (N=119)	SAVR (N=108)	TAVR (N=0)	SAVR (N=0)	TAVR (N=10)	SAVR (N=4)
n	1	2	90	86	0	0	7	3
Mean ± SD	0.57 ±	0.60 ± 0.00	0.72 ± 0.13	0.76 ± 0.12	NA	NA	0.66 ± 0.12	0.51 ± 0.20

	NYHA 1-2				NYHA 3-4			
	Major/Minor Stroke		No Stroke Before		Major/Minor Stroke		No Stroke Before	
	Before 6 Month		6 Month		Before 6 Month		6 Month	
	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR
SF-6D	(N=3)	(N=5)	(N=119)	(N=108)	(N=0)	(N=0)	(N=10)	(N=4)
Median	0.57	0.60	0.72	0.74	NA	NA	0.62	0.40
Min, Max	0.6, 0.6	0.6, 0.6	0.3, 1.0	0.5, 1.0	NA	NA	0.5, 0.8	0.4, 0.7
Q1, Q3	0.57, 0.57	0.60, 0.61	0.64, 0.80	0.67, 0.85	NA	NA	0.56, 0.81	0.39, 0.74

Supplementary Table 4: SF-6D and NYHA at 12-Month Visit

	NYHA 1-2				NYHA 3-4			
	Major/Minor Stroke		No Stroke Before		Major/Minor Stroke		No Stroke Before	
	Before 12 Month		12 Month		Before 12 Month		12 Month	
	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR
SF-6D	(N=3)	(N=5)	(N=125)	(N=111)	(N=0)	(N=0)	(N=4)	(N=4)
N	3	3	102	90	0	0	3	4

	NYHA 1-2				NYHA 3-4			
	Major/Minor Stroke		No Stroke Before		Major/Minor Stroke		No Stroke Before	
	Before 12 Month		12 Month		Before 12 Month		12 Month	
	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR
SF-6D	(N=3)	(N=5)	(N=125)	(N=111)	(N=0)	(N=0)	(N=4)	(N=4)
Mean ± SD	0.63 ± 0.04	0.69 ± 0.10	0.71 ± 0.12	0.75 ± 0.14	NA	NA	0.61 ± 0.12	0.57 ± 0.12
Median	0.63	0.66	0.73	0.74	NA	NA	0.63	0.53
Min, Max	0.6, 0.7	0.6, 0.8	0.3, 1.0	0.4, 1.0	NA	NA	0.5, 0.7	0.5, 0.7
Q1, Q3	0.60, 0.67	0.62, 0.80	0.64, 0.79	0.66, 0.85	NA	NA	0.47, 0.72	0.48, 0.65

Copyright

Supplementary Table 5: SF-6D and NYHA at 24-Month Visit


	NYHA 1-2				NYHA 3-4			
	Major/Minor Stroke		No Stroke Before		Major/Minor Stroke		No Stroke Before	
	Before 24 Month		24 Month		Before 24 Month		24 Month	
	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR	TAVR	SAVR
SF-6D	(N=4)	(N=5)	(N=115)	(N=105)	(N=0)	(N=0)	(N=4)	(N=4)
n	1	4	83	79	0	0	4	4
Mean ± SD	0.62 ±	0.66 ± 0.25	0.72 ± 0.12	0.73 ± 0.14	NA	NA	0.63 ± 0.17	0.60 ± 0.05
Median	0.62	0.61	0.73	0.74	NA	NA	0.68	0.59
Min, Max	0.6, 0.6	0.4, 1.0	0.4, 1.0	0.3, 1.0	NA	NA	0.4, 0.8	0.6, 0.7
Q1, Q3	0.62, 0.62	0.51, 0.81	0.66, 0.81	0.62, 0.84	NA	NA	0.51, 0.76	0.56, 0.64

Copy


Supplementary Table 6: Results of the Base Case and other Long-term Mortality Scenarios

	Costs (DKK)	Incremental Costs (DKK)	QALYs	Incremental QALYs	ICER (DKK/QALY)
<b>Fade-out Scenario I (proposed base case – fade out over 24 months to life tables [RR=1])</b>					
SAVR	211,581		5.30		
TAVI	276,142	64,561	5.39	0.09	696,264
<b>Fade-out Scenario II (fade out over 24 months to weighted RR [1.10] – then continues lifelong)</b>					
SAVR	210,826		5.19		
TAVI	275,329	64,503	5.28	0.09	712,188
<b>Continuous Benefit Scenario I (group-specific RRs [1.05/1.15] continue lifelong)</b>					
SAVR	210,484		5.14		
TAVI	275,722	64,238	5.33	0.19	334,166
<b>Stop-and-Drop Scenario I (immediate drop to life tables [RR=1.0])</b>					
SAVR	211,881		5.33		
TAVI	276,238	64,357	5.40	0.07	892,000
<b>Stop-and-Drop Scenario II (immediate drop to weighted RR [1.10] – but continues lifelong)</b>					
SAVR	210,920		5.20		
TAVI	275,239	64,319	5.27	0.07	904,070


## Supplementary Figures




Supplementary Figure 1: 60-month Mortality Data with Calibrated Model Output



Supplementary Figure 2: 60-month Stroke Incidence with Calibrated Model Output




Supplementary Figure 3: 60-Month Myocardial Infarction Incidence with Calibrated Model Output




Supplementary Figure 4: Tornado Diagram

### Incremental Cost-Effectiveness, TAVI v. SAVR




Supplementary Figure 5: Incremental Cost-Effectiveness (ICE) Scatter PLOT with a Willingness-to-Pay Threshold of DKK 375,489

### Incremental Cost-Effectiveness, TAVI v. SAVR




Supplementary Figure 6: Incremental Cost-Effectiveness (ICE) Scatter PLOT with a Willingness-to-Pay Threshold of DKK 1,126,467


**Disclaimer :** As a public service to our readership, this article -- peer reviewed by the Editors of EuroIntervention - has been published immediately upon acceptance as it was received. The content of this article is the sole responsibility of the authors, and not that of the journal




Supplementary Figure 7: Cost-effectiveness Acceptability Curve.




Supplementary Figure 8: Illustration of the Relative Mortality Risk Fading Out Over Time




Supplementary Figure 9: One-way Sensitivity Analyses of Relative Mortality Risk after TAVI and SAVR in the Fade-Out I Scenario (Base Case, Outcome: Undiscounted Life Years)




Supplementary Figure 10: Two-way Sensitivity Analysis on Relative Mortality Risk after SAVR over Relative Mortality Risk after TAVI (Fade-out Scenario)



Supplementary Figure 11: One-way Sensitivity Analyses of Relative Mortality Risk after TAVI and SAVR in the Continuous Benefit Scenario (Outcome: Undiscounted Life Years)




Supplementary Figure 12: One-way Sensitivity Analysis over Relative Mortality Risk after TAVI (Continuous Benefit Scenario)



Supplementary Figure 13: One-way Sensitivity Analysis Over Relative Mortality Risk after SAVR (Continuous Benefit Scenario)


Outcome: TAVI is highly cost-effective




Outcome: TAVI is cost-effective (but not highly cost-effective)

Outcome: TAVI is absolutely dominated by SAVR

Supplementary Figure 14: Two-way Sensitivity Analysis on Relative Mortality Risk after SAVR over Relative Mortality Risk after TAVI (Continuous Benefit Scenario)



Supplementary Figure 15: Bar Chart Comparing the QALYs Incurred by TAVI and SAVR



Supplementary Figure 16: Combined Bar and Line Chart Comparing the Incremental QALYs (blue bars) and corresponding ICERs (orange line)